

A bioinformatic approach to establish P38 α MAPK inhibitory mechanism of selected natural products in psoriasis

Anurag Agrawal^{1,2}, Rajendra Awasthi³, Giriraj T Kulkarni^{4*} & Lakshmayya⁵

¹School of Pharmacy, ITM University, Gwalior-474 001, Madhya Pradesh, India

²Institute of Pharmacy, Uttarakhand Technical University, Dehradun-248 007, Uttarakhand, India

³Amity Institute of Pharmacy, Amity University, Noida-201 303, Uttar Pradesh, India

⁴Gokaraju Rangaraju College of Pharmacy, Hyderabad-500 090, Telangana, India

⁵GRD-Institute of Management and Technology, Rajpur, Dehradun-248 009, Uttarakhand, India

Received 03 November 2020; revised 21 January 2022

Supplementary Data




Fig. S1 — Structures of ligands




Fig. S2 — 2D images of Interactions between natural ligands and Target enzyme (3lhj) (A) Caffeine; (B) Capsaicin; (C) Catechin; (D) Curcumin; (E) DMF; (F) Embelin; (G) Epicatechin; (H) Gossypol; (I) Hypericin; (J) Luteolin; (K) MEF; and (L) Quercetin (Contd.)




Fig. S2 — 2D images of Interactions between natural ligands and Target enzyme (3lhj) (A) Caffeine; (B) Capsaicin; (C) Catechin; (D) Curcumin; (E) DMF; (F) Embelin; (G) Epicatechin; (H) Gossypol; (I) Hypericin; (J) Luteolin; (K) MEF; and (L) Quercetin (*Contd.*)