

substitution by the methylene group at C-8 carbon in one chromanone moiety along with two singlets at δ 6.42 and 7.90 (each integrating for one proton) indicating substitution by the methylene group at C-6 carbon of the other chromanone moiety. Hence, **9** was characterized as the unsymmetrical 6,8'-isomer. Comparison of the spectral data of **10** with that of **8** and **9** indicated it to be another of their symmetrical methylene *bis* isomer. Compound **10** formed as a minor product, displayed only two singlets at δ 6.43 and 7.85 (integrating for two protons each), in the aromatic region of its ^1H NMR spectrum, assigned to H-8 and 8' and H-5 and 5' protons, respectively, thus confirming its structure.

O-Methoxyacetyl derivative **2** produced isomeric methylene-*bis*-compounds **11** (in minor amount) and **12**. These were identified by detailed spectral studies on a similar basis as for **8**, **9** and **10**. The ^1H NMR spectrum of **11** showed the presence of only one pair of *ortho*-coupled doublets at δ 6.59 and 6.99 (each integrating for two protons) indicating substitution at C-5 carbon in both chromanone moieties and hence was identified as the symmetrical 5,5'-isomer. Comparison of spectral data of **12** with that of **11** indicated it to be its unsymmetrical isomer. In the aromatic region of its ^1H NMR spectrum, the presence of a pair of *ortho*-coupled doublets at δ 6.87 and 7.15 (each integrating for one proton) indicated substitution at C-5 carbon in one chromanone moiety while two singlets at δ 7.01 and 7.34 (each integrating for one proton) indicated substitution at C-7 carbon in the other chromanone moiety, thereby confirming it to be the 5,7'-isomer (Scheme I).

O-Methoxyacetyl derivative **3** upon standard condition treatment furnished furanocoumarin¹² **13** and methylene *bis* product **14** which was identified by detailed spectral analysis as the symmetrical 6,6'-isomer. Similarly, **4** and **5** produced furanocoumarins¹² **15** and **18** and isomeric methylene-*bis*-coumarins **16**, **17** and **19**, **20**, respectively, which were characterized by detailed spectral analysis on similar basis as discussed above (Scheme III).


O-Methoxyacetyl derivative **6** upon reaction gave **21** (in minor amount) and **22** while **7** gave **23** and **24** as the major products along with minor amount of **25** (Scheme II). These were also characterized by detailed spectral studies similar to the discussion above.

For the purpose of synthesising methylene *bis* compounds using Fries reaction starting from 3- and 4-hydroxycoumarins **26** and **28**, the corresponding *O*-

methoxyacetyl derivatives were obtained in poor yields and hence, further reaction could not be done. However, when subjected to Friedel-Crafts reaction with methoxyacetyl chloride in presence of anhydrous AlCl_3 in CS_2 solvent, **26** and **28** afforded corresponding methylene *bis* coumarins **27** and **29**, respectively (Scheme IV). Their ^1H NMR indicated no substitution in the benzene ring and absence of H attached directly to the pyrone ring. Detailed study of their spectral data was used to characterize them. The presence of methylene bridge in pyrone ring of **27** and **29** was at position *ortho* to the hydroxy group and was in agreement with the products formed in Scheme I, Scheme II and Scheme III.

To explain this novel product formation in the reaction, we propose a mechanism which involves generation of an unstable methoxyacyl cation that upon loss of CO generates a resonance stabilized methoxymethyl cation **30**, which subsequently undergoes electrophilic substitution on the phenolic nucleus of the benzopyrone moiety to form corresponding benzyl methyl ether **31**. Subsequent complexation with a Lewis acid generates the corresponding benzyl cation which on electrophilic substitution on another phenolic nucleus of the other benzopyrone moiety forms the methylene *bis* compound **33** (Scheme V).

The structures of all the synthesized compounds, *O*-methoxyacetyl derivatives (**1-7**), methylene-*bis*-compounds and furanocoumarins (**8-29**) were unambiguously established on the basis of their spectral (IR, ^1H -, ^{13}C -NMR, ^1H - ^1H COSY NMR, ^1H - ^{13}C HMBC NMR and HRMS) data analysis. The structure of known compounds **13**, **15** and **18** were further confirmed by the comparison of their physical and spectral data with those reported in literature¹².



Scheme V — Plausible mechanism for the synthesis of methylene-bis-compounds

Experimental Section

All chemicals were purchased from Sigma-Aldrich Chemicals Pvt. Limited India and from local commercial sources. Solvents were removed under reduced pressure using rotary evaporator, followed by further removal of the residual solvent under high vacuum. Column chromatography was performed over silica gel (60-120 mesh). Melting points were determined on Buchi M-560 instrument and are uncorrected. HRMS analysis was carried out using Agilent G6530AA LC Q-TOF mass spectrometer using ESI method. The IR spectra of compounds were recorded on Perkin-Elmer model 2000 FT-IR spectrometer and are expressed as wavenumber (cm^{-1}). The ^1H - and the ^{13}C - and other 2D NMR spectra were recorded on 250 or 300 MHz spectrometer using tetramethylsilane (TMS) as internal standard. The chemical shift values are on δ scale and the coupling constant (J) are in Hz.

General procedure for the synthesis of *O*-methoxyacetyl derivatives, 1-7

The corresponding hydroxyl compounds of 1-7 were prepared by earlier known procedures¹³⁻¹⁵. Dry pyridine was added drop-wise to a stirred solution of each of these hydroxy benzopyrones (10 mmol) followed by addition of methoxyacetyl chloride (15 mmol) under anhydrous conditions. The reaction mixture was warmed to 60-70°C on a water bath for 30 min and then left at room temperature for 24 h. The contents were then poured over an ice-water mixture and the precipitate thus formed was collected by filtration, washed with water, dried and crystallized from DCM-MeOH to afford compounds 1-7 in 90-98% yields.

Note: The hydroxy chromanones were prepared by catalytic hydrogenation of the corresponding chromones.

7-*O*-Methoxyacetylchromanone, 1: Colourless needles (94%); m.p. 117-118°C; IR cm^{-1} (KBr): 1765, 1653; ^1H NMR (300 MHz, CDCl_3): δ 2.71 (t, $J = 6.4$

Hz, 2H, H-3), 3.55 (s, 3H, -OCH₃), 4.34 (s, 2H, -OCH₂-), 4.48 (t, $J = 6.6$ Hz, 2H, H-2), 6.35 (d, $J = 2.3$ Hz, 1H, H-8), 6.54 (dd, $J = 2.3$ and 8.7 Hz, 1H, H-6), 7.77 (d, $J = 8.7$ Hz, 1H, H-5); ^{13}C NMR (75.5 MHz, CDCl_3): δ 36.8, 60.1, 67.3, 69.2, 101.1, 109.9, 115.2, 129.0, 163.8, 165.8, 168.8, 191.1; MS m/z 236 (M^+). Anal. Calcd for $\text{C}_{12}\text{H}_{12}\text{O}_5$: C, 60.99; H, 5.12. Found: C, 60.93; H, 5.11.

6-*O*-Methoxyacetylchromanone, 2: Colourless needles (95%); m.p.: 123°C; IR cm^{-1} (KBr): 1763, 1646; ^1H NMR (300 MHz, CDCl_3): δ 2.77 (t, $J = 6.6$ Hz, 2H, H-3), 3.56 (s, 3H, -OCH₃), 4.34 (s, 2H, -OCH₂-), 4.48 (t, $J = 6.6$ Hz, 2H, H-2), 6.87 (d, $J = 8.7$ Hz, 1H, H-8), 7.02 (dd, $J = 2.1$ and 8.7 Hz, 1H, H-7), 7.28 (d, $J = 2.2$ Hz, 1H, H-5); ^{13}C NMR (75.5 MHz, CDCl_3): δ 37.5, 59.8, 67.2, 68.9, 108.1, 119.1, 120.9, 125.6, 150.3, 158.1, 168.7, 192.0; MS m/z 236 (M^+). Anal. Calcd for $\text{C}_{12}\text{H}_{12}\text{O}_5$: C, 60.99; H, 5.12. Found: C, 61.07; H, 5.16.

5-*O*-Methoxyacetyl-4-methyl coumarin, 3: Colourless needles (91%); m.p.: 125-127°C; UV nm (MeOH): 316, 268, 227; IR cm^{-1} (KBr): 1766, 1697; ^1H NMR (250 MHz, CDCl_3): δ 2.43 (d, $J = 1.0$ Hz, 3H, -CH₃ at C-4), 3.58 (s, 3H, -OCH₃), 4.34 (s, 2H, -OCH₂-), 6.35 (q, $J = 1.0$ Hz, 1H, H-3), 7.29-7.41 (m, 3H, H-6, 7 and 8); ^{13}C NMR (62.9 MHz, CDCl_3): δ 18.5, 59.6, 69.8, 115.9, 117.0, 118.1, 120.7, 125.0, 146.2, 151.2, 151.5, 160.2, 168.7; MS m/z 248 (M^+). Anal. Calcd for $\text{C}_{13}\text{H}_{12}\text{O}_5$: C, 62.88; H, 4.87. Found: C, 62.92; H, 4.86.

6-*O*-Methoxyacetyl-4-methyl coumarin, 4: Colourless needles (93%); m.p.: 117-118°C; UV nm (MeOH): 318, 269, 224; IR cm^{-1} (KBr): 1770, 1720; ^1H NMR (250 MHz, CDCl_3): δ 2.42 (d, $J = 1.1$ Hz, 3H, -CH₃ at C-4), 3.56 (s, 3H, -OCH₃), 4.33 (s, 2H, -OCH₂-), 6.33 (q, $J = 1.1$ Hz, 1H, H-3), 7.31 (d, $J = 8.8$ Hz, 1H, H-8), 7.34 (d, $J = 1.9$ Hz, 1H, H-5), 7.38 (dd, $J = 1.9$ and 8.8 Hz, 1H, H-7); ^{13}C NMR (62.9 MHz, CDCl_3): δ 18.6, 59.6, 69.7, 115.8, 117.1, 118.1, 125.0, 130.1, 146.1, 151.1, 151.7, 160.3, 168.8; MS

m/z 248 (M^+). Anal. Calcd for $C_{13}H_{12}O_5$: C, 62.88; H, 4.87. Found: C, 62.89; H, 4.89.

7-*O*-Methoxyacetyl-4-methyl coumarin, 5: Colourless needles (92%); m.p.: 136°C; UV nm (MeOH): 317, 270, 225; IR cm^{-1} (KBr): 1763, 1715; 1H NMR (250 MHz, $CDCl_3$): δ 2.42 (d, $J = 1.0$ Hz, 3H, -CH₃ at C-4), 3.54 (s, 3H, -OCH₃), 4.32 (s, 2H, -OCH₂-), 6.26 (q, $J = 1.0$ Hz, 1H, H-3), 7.11 (dd, $J = 2.1$ and 8.8 Hz, 1H, H-6), 7.14 (d, $J = 2.1$ Hz, 1H, H-8), 7.64 (d, $J = 8.8$ Hz, 1H, H-7); ^{13}C NMR (62.9 MHz, $CDCl_3$): δ 18.6, 59.5, 69.4, 110.1, 114.5, 117.7, 117.9, 125.4, 151.7, 152.2, 153.9, 160.2, 168.05; MS m/z 248 (M^+). Anal. Calcd for $C_{13}H_{12}O_5$: C, 62.88; H, 4.87. Found: C, 62.78; H, 4.84.

6-*O*-Methoxyacetylchromone, 6: Colourless needles (94%); m.p.: 176°C; IR cm^{-1} (KBr): 1765, 1638; 1H NMR (300 MHz, $CDCl_3$): δ 3.55 (s, 3H, -OCH₃), 4.31 (s, 2H, -OCH₂-), 6.35 (d, $J = 6.0$ Hz, 1H, H-3), 7.45 (dd, $J = 2.6$ and 9.0 Hz, 1H, H-7), 7.52 (d, $J = 9.0$ Hz, 1H, H-8), 7.87 (d, $J = 6.0$ Hz, 1H, H-2), 7.93 (d, $J = 2.6$ Hz, 1H, H-5); ^{13}C NMR (75.5 MHz, $CDCl_3$): δ 59.6, 69.6, 112.6, 117.8, 119.8, 125.7, 127.8, 147.1, 154.2, 155.5, 168.7, 184.8; MS m/z 234 (M^+). Anal. Calcd for $C_{12}H_{10}O_5$: C, 61.52; H, 4.31. Found: C, 61.47; H, 4.29.

7-*O*-Methoxyacetylchromone, 7: Colourless needles (97%); m.p.: 187-188°C; IR cm^{-1} (KBr): 1762, 1635; 1H NMR (300 MHz, $CDCl_3$): δ 3.61 (s, 3H, -OCH₃), 4.42 (s, 2H, -OCH₂-), 6.36 (d, $J = 5.9$ Hz, 1H, H-3), 6.80 (d, $J = 2.4$ Hz, 1H, H-8), 6.94 (dd, $J = 2.4$ and 8.8 Hz, 1H, H-6), 7.87 (d, $J = 5.9$ Hz, 1H, H-2), 7.98 (d, $J = 8.8$ Hz, 1H, H-5); ^{13}C NMR (75.5 MHz, $CDCl_3$): δ 58.9, 68.8, 103.0, 111.1, 114.6, 116.1, 130.5, 155.2, 164.1, 166.9, 168.7, 169.1, 185.2; MS m/z 234 (M^+). Anal. Calcd for $C_{12}H_{10}O_5$: C, 61.52; H, 4.31. Found: C, 61.50; H, 4.27.

General procedure for Lewis acid mediated rearrangement of *O*-methoxyacetyl derivatives, 1-7

A mixture of *O*-methoxyacetyl derivative (1-7, 10 mmol) and anhydrous $AlCl_3$ (25 mmol) was heated slowly to 170°C over a period of 4 h. The complex formed was then decomposed by adding it to an ice-water mixture and acidifying the resulting solution with HCl (10 ml). The contents were heated on a water bath for 30 min, cooled and the solid product formed was collected, washed with water, dried and purified by column chromatography to afford methylene *bis*-compounds 8-25 in 60 to 80% yields.

8,8'-Methylene-bis(7-hydroxychromanone), 8: Elution with $CHCl_3/MeOH$ (97:3) gave 8(40%); m.p.

> 300°C; UV nm (MeOH): 319, 278, 233; IR cm^{-1} (KBr): 3279, 1647; 1H NMR (300 MHz, $DMSO-d_6$): δ 2.50 (t, $J = 6.3$ Hz, 4H, H-3 and 3'), 3.68 (s, 2H, -CH₂-), 4.27 (t, $J = 6.3$ Hz, 4H, H-2 and 2'), 6.49 (d, $J = 8.6$ Hz, 2H, H-6 and 6'), 7.35 (d, $J = 8.6$ Hz, 2H, H-5 and 5'), 9.98 (brs, 2H, 2 \times -OH); ^{13}C NMR (75.5 MHz, $DMSO-d_6$): δ 17.2, 37.3, 67.3, 109.9, 114.3, 114.5, 125.6, 162.1, 162.4, 190.8; MS m/z 340 (M^+). Anal. Calcd for $C_{19}H_{16}O_6$: C, 67.04; H, 4.74. Found: C, 67.10; H, 4.79.

6,8'-Methylene-bis(7-hydroxychromanone), 9: Elution with $CHCl_3/MeOH$ (97:3) gave 9 in the earlier fractions followed by 9 in the later fractions (28%); m.p. > 300°C; UV nm (MeOH): 320, 247, 212; IR cm^{-1} (KBr): 3435, 1649; 1H NMR (300 MHz, $DMSO-d_6$): δ 2.61 and 2.68 (t, $J = 6.4$ Hz, 2H each, H-3 and 3'), 3.29 (s, 2H, -CH₂-), 4.42 and 4.45 (t, $J = 6.4$ Hz, 2H each, H-2 and 2'), 6.42 (s, 1H, H-8), 6.65 (d, $J = 8.6$ Hz, 1H, H-6'), 7.65 (d, $J = 8.6$ Hz, 1H, H-5'), 7.90 (s, 1H, H-5), 10.04 (brs, 2H, 2 \times -OH); ^{13}C NMR (75.5 MHz, $DMSO-d_6$): δ 21.6, 37.0, 37.2, 66.9, 67.1, 102.1, 109.9, 113.1, 113.5, 121.7, 126.4, 161.8, 162.6, 190.3; MS m/z 340 (M^+). Anal. Calcd for $C_{19}H_{16}O_6$: C, 67.04; H, 4.74. Found: C, 67.02; H, 4.77.

6,6'-Methylene-bis(7-hydroxychromanone), 10: Elution with $CHCl_3/MeOH$ (95:5) gave 10 (12%); m.p. > 300°C; UV nm (MeOH): 319, 247, 211; IR cm^{-1} (KBr): 3395, 1651; 1H NMR (300 MHz, $DMSO-d_6$): δ 2.57 (t, $J = 6.4$ Hz, 4H, H-3 and 3'), 3.51 (s, 2H, -CH₂-), 4.31 (t, $J = 6.4$ Hz, 4H, H-2 and 2'), 6.43 (s, 2H, H-8 and 8'), 7.85 (s, 2H, H-5 and 5'), 10.87 (brs, 2H, 2 \times -OH); ^{13}C NMR (75.5 MHz, $DMSO-d_6$): δ 22.1, 37.3, 67.2, 102.7, 114.1, 120.9, 125.7, 161.9, 162.5, 190.7; MS m/z 340 (M^+). Anal. Calcd for $C_{19}H_{16}O_6$: C, 67.04; H, 4.74. Found: C, 67.09; H, 4.75.

5,5'-Methylene-bis(6-hydroxychromanone), 11: Elution with $CHCl_3/MeOH$ (95:5) gave 11 (18%); m.p. > 300°C; UV nm (MeOH): 321, 249, 210; IR cm^{-1} (KBr): 3391, 1645; 1H NMR (300 MHz, $DMSO-d_6$): δ 2.56 (t, $J = 6.4$ Hz, 4H, H-3 and 3'), 3.67 (s, 2H, -CH₂-), 4.24 (t, $J = 6.4$ Hz, 4H, H-2 and 2'), 6.59 (d, $J = 8.6$ Hz, 2H, H-8 and 8'), 6.99 (d, $J = 8.6$ Hz, 2H, H-7 and 7'), 10.03 (brs, 2H, 2 \times -OH); ^{13}C NMR (75.5 MHz, $DMSO-d_6$): δ 18.2, 37.6, 67.4, 105.7, 119.9, 120.1, 137.9, 147.1, 157.0, 191.9; MS m/z 340 (M^+). Anal. Calcd for $C_{19}H_{16}O_6$: C, 67.04; H, 4.74. Found: C, 67.08; H, 4.75.

5,7'-Methylene-bis(6-hydroxychromanone), 12: Elution with $CHCl_3/MeOH$ (93:7) gave 12 (55%);

m.p. > 300°C; UV nm (MeOH): 321, 250, 211; IR cm^{-1} (KBr): 3382, 1637; ^1H NMR (300 MHz, DMSO- d_6): δ 2.66 and 2.73 (t, $J = 6.5$ Hz, 2H each, H-3 and 3'), 3.34 (s, 2H, $-\text{CH}_2-$), 4.43 and 4.46 (t, $J = 6.5$ Hz, 2H each, H-2 and 2'), 6.87 (d, $J = 8.6$ Hz, 1H, H-8), 7.01 (s, 1H, H-8'), 7.15 (d, $J = 8.6$ Hz, 1H, H-7), 7.34 (s, 1H, H-5'), 9.95 (brs, 2H, 2 \times -OH); ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 21.1, 37.4, 37.5, 67.0, 67.1, 106.4, 118.7, 120.0, 127.7, 131.4, 137.5, 146.9, 156.7, 190.8, 191.4; MS m/z 340 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{16}\text{O}_6$: C, 67.04; H, 4.74. Found: C, 66.98; H, 4.69.

Furan-2(3H)-one[5,4-f]-4-methyl-2H-1-benzopyran-2-one, 13: All spectral data are identical to the reported data¹².

6,6'-Methylene-bis(5-hydroxy-4-methylcoumarin), 14: Elution with $\text{CHCl}_3/\text{MeOH}$ (92:8) gave **14** (62%); m.p.>300°C; UV nm (MeOH): 295, 255, 210; IR cm^{-1} (KBr): 3300, 1690; ^1H NMR (250 MHz, DMSO- d_6): δ 2.46 (d, $J = 1.1$ Hz, 6H, 2 \times - CH_3), 4.21 (s, 2H, $-\text{CH}_2-$), 6.32 (q, $J = 1.1$ Hz, 2H, H-3 and 3'), 6.93 (d, $J = 8.6$ Hz, 2H, H-8 and 8'), 7.45 (d, $J = 8.6$ Hz, 2H, H-7 and 7'), 9.79 (brs, 2H, 2 \times -OH); ^{13}C NMR (62.9 MHz, DMSO- d_6): δ 18.4, 27.9, 113.8, 116.4, 117.5, 121.9, 126.7, 146.3, 151.4, 155.1, 159.4; MS m/z 364 (M^+). Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{O}_6$: C, 69.21; H, 4.44. Found: C, 69.15; H, 4.39.

Furan-2(3H)-one[4,5-f]-4-methyl-2H-1-benzopyran-2-one, 15: All spectral data are identical to the reported data¹².

5,7'-Methylene-bis(6-hydroxy-4-methylcoumarin), 16: Elution with $\text{CHCl}_3/\text{MeOH}$ (95:5) gave **16** (18%); m.p.>300°C; UV nm (MeOH): 346, 280, 233; IR cm^{-1} (KBr): 3182, 1674; ^1H NMR (250 MHz, DMSO- d_6): δ 2.35 (d, $J = 1.0$ Hz, 3H, $-\text{CH}_3$), 2.36 (d, $J = 1.0$ Hz, 3H, $-\text{CH}_3$), 4.33 (s, 2H, $-\text{CH}_2-$), 6.26 and 6.28 (s, 1H each, H-3 and 3'), 7.12 (s, 1H, H-8'), 7.15 (d, $J = 8.8$ Hz, 1H, H-8), 7.27 (d, $J = 8.8$ Hz, 1H, H-7), 7.28 (s, 1H, H-5'), 9.82 and 10.06 (brs, 1H each, 2 \times -OH); ^{13}C NMR (62.9 MHz, DMSO- d_6): δ 17.9, 22.7, 26.9, 108.5, 109.4, 116.7, 17.8, 119.0, 119.6, 121.2, 130.0, 133.2, 140.6, 146.3, 147.8, 151.1, 152.5, 152.7, 153.6, 159.3, 159.8; MS m/z 364 (M^+). Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{O}_6$: C, 69.21; H, 4.44. Found: C, 69.22; H, 4.41.

7,7'-Methylene-bis(6-hydroxy-4-methylcoumarin), 17: Elution with $\text{CHCl}_3/\text{MeOH}$ (90:10) gave **17** (42%); m.p.>300°C; UV nm (MeOH): 343, 280, 226; IR cm^{-1} (KBr): 3221, 1686; ^1H NMR (250 MHz,

DMSO- d_6): δ 2.50 (d, $J = 1.5$ Hz, 6H, 2 \times - CH_3), 3.97 (s, 2H, $-\text{CH}_2-$), 6.32 (m, 2H, H-3 and 3'), 7.04 (s, 2H, H-8 and 8'), 7.21 (s, 2H, H-5 and 5'), 9.98 (brs, 2H each, 2 \times -OH); ^{13}C NMR (62.89 MHz, DMSO- d_6): δ 18.4, 28.8, 110.4, 117.3, 119.7, 131.1, 133.1, 146.8, 151.9, 153.0, 160.5; MS m/z 364 (M^+). Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{O}_6$: C, 69.21; H, 4.44. Found: C, 69.25; H, 4.49.

Furan-2(3H)-one[5,4-h]-4-methyl-2H-1-benzopyran-2-one, 18: All spectral data are identical to the reported data¹².

8,8'-Methylene-bis(7-hydroxy-4-methylcoumarin), 19: Elution with $\text{CHCl}_3/\text{MeOH}$ (96:4) gave **19** (48%); m.p.>300°C; UV nm (MeOH): 330, 230; IR cm^{-1} (KBr): 3320, 1690; ^1H NMR (250 MHz, DMSO- d_6): δ 2.42 (s, 6H, 2 \times - CH_3), 4.25 (s, 2H, $-\text{CH}_2-$), 6.15 (s, 2H, H-3 and 3'), 6.83 (d, $J = 9$ Hz, 2H, H-6 and 6'), 7.50 (d, $J = 9$ Hz, 2H, H-5 and 5'), 10.31 (brs, 2H, 2 \times -OH); ^{13}C NMR (62.89 MHz, DMSO- d_6): δ 17.3, 18.6, 110.2, 113.6, 123.9, 153.4, 154.2, 159.3, 160.9; MS m/z 364 (M^+). Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{O}_6$: C, 69.21; H, 4.44. Found: C, 70.17; H, 4.57.

6,8'-Methylene-bis(7-hydroxy-4-methylcoumarin), 20: Elution with $\text{CHCl}_3/\text{MeOH}$ (92:8) gave **20** (22%); m.p.>300°C; UV nm (MeOH): 331, 232; IR cm^{-1} (KBr): 3330, 1700; ^1H NMR (250 MHz, DMSO- d_6): δ 2.36 (s, 6H, 2 \times - CH_3), 3.86 (s, 2H, $-\text{CH}_2-$), 6.09 (s, 2H, H-3 and 3'), 6.81 (d, $J = 9$ Hz, 1H, H-6'), 6.78 (s, 1H, H-8), 7.26 (s, 1H, H-5), 7.64 (d, $J = 9$ Hz, H-5') 10.45 and 10.79 (s, 1H each, 2 \times -OH); ^{13}C NMR (62.9 MHz, DMSO- d_6): δ 18.1, 21.7, 26.4, 107.5, 108.3, 116.1, 117.2, 118.8, 119.1, 121.3, 130.1, 132.9, 141.1, 146.2, 147.6, 151.4, 152.5, 152.9, 153.6, 159.3, 159.7; MS m/z 364 (M^+). Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{O}_6$: C, 69.21; H, 4.44. Found: C, 68.92; H, 4.50.

5,5'-Methylene-bis(6-hydroxychromone), 21: Elution with $\text{CHCl}_3/\text{MeOH}$ (95:5) gave **21** (15%); m.p.>300°C; UV nm (MeOH): 310, 248, 212 nm; IR cm^{-1} (KBr): 3089, 1631; ^1H NMR (300 MHz, DMSO- d_6): δ 4.08 (s, 2H, $-\text{CH}_2-$), 6.19 (d, $J = 5.9$ Hz, 2H, H-2 and 2'), 7.10 (d, $J = 9$ Hz, 2H, H-8 and 8'), 7.21 (d, $J = 9$ Hz, 2H, H-7 and 7'), 7.79 (d, $J = 5.9$ Hz, 2H, H-2 and 2') 10.48 (brs, 2H each, 2 \times -OH); ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 18.3, 110.2, 117.5, 121.1, 129.5, 137.0, 146.2, 154.2, 154.4, 184.8; MS m/z 336 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{12}\text{O}_6$: C, 67.84; H, 3.59. Found: C, 67.79; H, 3.57.

5,7'-Methylene-bis(6-hydroxychromone), 22: Elution with $\text{CHCl}_3/\text{MeOH}$ (93:7) gave **22** (52%);

m.p. > 300°C; UV nm (MeOH): 310, 247, 211 nm; IR cm^{-1} (KBr): 3098, 1629; ^1H NMR (300 MHz, DMSO- d_6): δ 4.14 (s, 2H, $-\text{CH}_2-$), 6.24 and 6.29 (d, $J = 6.0$ Hz, 1H each, H-3 and 3'), 7.28 (d, $J = 9.1$ Hz, 1H, H-7), 7.35 (d, $J = 9.1$ Hz, 1H, H-8), 7.45 (s, 1H, H-8'), 7.59 (s, 1H, H-5'), 7.85 and 7.92 (d, $J = 6.0$ Hz, 1H each, H-2 and 2'), 10.61 (brs, 2H each, 2 \times -OH); ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 17.9, 110.5, 117.3, 117.4, 121.5, 126.8, 129.5, 136.1, 138.9, 146.8, 154.2, 154.3, 154.5, 184.9; MS m/z 336 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{12}\text{O}_6$: C, 67.84; H, 3.59. Found: C, 67.82; H, 3.60.

6,8'-Methylene-bis(7-hydroxychromone), 23: Elution with $\text{CHCl}_3/\text{MeOH}$ (94:6) gave **23** (26%); m.p. > 300°C; UV nm (MeOH): 306, 252, 211 nm; IR cm^{-1} (KBr): 3098, 1628; ^1H NMR (300 MHz, DMSO- d_6): δ 4.12 (s, 2H, $-\text{CH}_2-$), 6.10 and 6.17 (d, $J = 5.9$ Hz, 1H each, H-3 and 3'), 7.07 (d, $J = 8.7$ Hz, 1H, H-6'), 7.17 (s, 1H, H-8), 7.89 (d, $J = 8.7$ Hz, 1H, H-5'), 7.99 (s, 1H, H-5), 8.03 and 8.07 (d, $J = 5.9$ Hz, 1H each, H-2 and 2'), 10.62 (brs, 2H each, 2 \times -OH); ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 17.7, 103.2, 111.0, 114.1, 114.2, 115.2, 115.9, 123.1, 130.2, 155.4, 155.8, 163.0, 164.0, 185.1, 185.2; MS m/z 336 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{12}\text{O}_6$: C, 67.84; H, 3.59. Found: C, 67.87; H, 3.64.

8,8'-Methylene-bis(7-hydroxychromone), 24: Elution with $\text{CHCl}_3/\text{MeOH}$ (94:6) in the later fractions gave **24** (42%); m.p. > 300°C; UV nm (MeOH): 307, 252, 211 nm; IR cm^{-1} (KBr): 3099, 1630; ^1H NMR (300 MHz, DMSO- d_6): δ 4.27 (s, 2H, $-\text{CH}_2-$), 6.09 (d, $J = 5.8$ Hz, 2H, H-3 and 3'), 6.83 (d, $J = 8.7$ Hz, 2H, H-6 and 6'), 7.70 (d, $J = 8.7$ Hz, 2H, H-5 and 5'), 7.98 (d, $J = 5.8$ Hz, 2H each, H-2 and 2'), 10.51 (brs, 2H each, 2 \times -OH); ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 17.5, 110.9, 114.1, 115.3, 116.1, 130.5, 155.6, 162.5, 163.9, 185.4; MS m/z 336 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{12}\text{O}_6$: C, 67.84; H, 3.59. Found: C, 67.83; H, 3.62.

6,6'-Methylene-bis(7-hydroxychromone), 25: Elution with $\text{CHCl}_3/\text{MeOH}$ (90:10) gave **25** (16%); m.p. > 300°C; UV nm (MeOH): 307, 252, 211; IR cm^{-1} (KBr): 3110, 1635; ^1H NMR (300 MHz, DMSO- d_6): δ 4.17 (s, 2H, $-\text{CH}_2-$), 6.14 (d, $J = 5.8$ Hz, 2H, H-3 and 3'), 6.98 (s, 2H, H-8 and 8'), 7.92 (s, 2H, H-5 and 5'), 7.96 (d, $J = 5.8$ Hz, 2H, H-2 and 2'), 10.73 (brs, 2H, 2 \times -OH); ^{13}C NMR (75.5 MHz, DMSO- d_6): δ 17.9, 103.2, 114.0, 115.9, 122.9, 130.3, 156.1, 162.8, 163.6, 184.9; MS m/z 336 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{12}\text{O}_6$: C, 67.84; H, 3.59. Found: C, 67.80; H, 3.57.

General procedure for the synthesis of methylene bis derivatives **27** and **29** from 3- and 4-hydroxycoumarin

A mixture of 3- or 4-hydroxycoumarin (10 mmol), anhydrous AlCl_3 (25 mmol) and freshly distilled CS_2 (15 mL) was stirred for 30 min under an inert atmosphere and methoxyacetyl chloride (15 mmol) was added to it. The contents were stirred for another 30 min at RT and then refluxed for 2 h. The apparatus was assembled for downward distillation and CS_2 was removed completely. The remaining contents were added to ice-water mixture and the solid that precipitated was collected by filtration and crystallised from ethanol to give a white solid of compound **27** and **29** respectively.

4,4'-Methylene-bis(3-hydroxycoumarin), 27: Yield 53%; m.p. 265-266°C; IR cm^{-1} (KBr): 3450, 1680; ^1H NMR (250 MHz, DMSO- d_6): δ 3.82 (s, 2H, $-\text{CH}_2-$), 6.56 (brs, 2H, 2 \times -OH), 7.30 (m, 2H), 7.32 (m, 2H), 7.35 (m, 4H); ^{13}C NMR (62.9 MHz, DMSO- d_6): δ 20.6, 114.6, 116.6, 127.6, 129.7, 131.6, 135.3, 140.4, 142.1, 159.7; MS m/z 336 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{12}\text{O}_6$: C, 67.84; H, 3.59. Found: C, 67.94; H, 3.63.

3,3'-Methylene-bis(4-hydroxycoumarin), 29: Yield 51%; m.p. 288°C; IR cm^{-1} (KBr): 3500, 1657; ^1H NMR (250 MHz, DMSO- d_6): δ 3.85 (s, 2H, $-\text{CH}_2-$), 7.37 (m, 2H), 7.39 (dd, 2H, $J = 2.0, 8.1$ Hz), m, 2H), 7.60 (m, 2H), 8.01 (dd, 2H, $J = 1.5, 8.0$ Hz), 11.33 (brs, 2H, 2 \times -OH); ^{13}C NMR (62.9 MHz, DMSO- d_6): δ 20.1, 102.7, 116.8, 120.2, 124.1, 124.6, 132.4, 153.2, 166.1, 169.7; MS m/z 336 (M^+). Anal. Calcd for $\text{C}_{19}\text{H}_{12}\text{O}_6$: C, 67.84; H, 3.59. Found: C, 67.89; H, 3.62.

Conclusion

An efficient Lewis acid promoted synthesis of methylene α - and γ -bis-benzopyrones was achieved wherein the linking of two benzopyrone moieties was *via* a methylene bridge on their benzene ring. The reaction proceeded without any solvent system and produced isomeric methylene bis derivatives of chromanones, coumarin and chromones in good yields. Earlier reported formation of furanocoumarins in case of Fries rearrangement on 5-, 6- and 7-methoxyacetyl coumarins due to intramolecular rearrangement was not observed in case of corresponding reactions on chromanones and chromones, which gave only isomeric methylene-bis-derivatives by an intermolecular rearrangement in good yields. Friedel-Crafts reaction on 3- and 4-

hydroxycoumarins led to the formation of methylene bis-derivatives.

Acknowledgement

The authors are grateful to Institute of Imminence, University of Delhi for providing financial support to strengthen research and development. The authors are thankful to DST-FIST program, CIF-USIC University of Delhi for providing the NMR spectral and HRMS recording facilities. Sandeep Kumar thanks CSIR, New Delhi for award of SPM Research Fellowship [File No. 09/045(0269)/2018-EMR-1].

References

- 1 Shahzad D, Saeed A, Faisal M, Larik F A, Bilquees S & Channar P A, *Org Prep Proced Int*, 51 (2019) 199.
- 2 Khan K M, Rahim F, Wadood A, Kosar N, Taha M, Lalani S, Khan A, Fakhri M I, Junaid M, Rehman W, Khan M, Perveen S, Sajid M & Choudhary M I, *Eur J Med Chem*, 81 (2014) 245.
- 3 Su C X, Mouscadet J F, Chiang C C, Tsai H J & Hsu L Y, *Chem Pharm Bull*, 54 (2006) 682.
- 4 Baruah P, Basumatary G, Yesylevskyy S O, Aguan K, Bez G & Mitra S, *J Biomol Str.Dynamics*, 37 (2018) 1750.
- 5 Reis J, Gaspar A, Milhazes N & Borges F, *J Med Chem*, 60 (2017) 7941.
- 6 Mayank, Singh A, Kaur N, Garg N & Singh N, *Comp Bio Chem*, 83 (2019) 107104.
- 7 Kostova I, Manolov I & Momekov G, *Eur J Med Chem*, 39 (2004) 765.
- 8 Mehrabi H & Abusaidi H, *J Iran Chem Soc*, 7 (2010) 890.
- 9 Heravi M M, Nahavandi F, Sadjadi S, Oskooie H A & Bamoharram F F, *Synth Commun*, 40 (2010) 498.
- 10 Gorbulevko N V, Tkachuk T M, Shokol T V, Semeniuchenko V V, Turov A V & Khilya V P, *Chem Heterocycl Comp*, 43 (2007) 683.
- 11 Nagaraj A & Reddy C S, *J Heterocycl Chem*, 44 (2007) 1357.
- 12 Babu B R, Bali S, Sakhuja R & Jain S C, *Indian J Chem*, 45B (2006) 2546.
- 13 Trivedi K N & Sethna S, *J Org Chem*, 25 (1960) 1817.
- 14 Sengupta P, Sen M, Karuri P, Wenkert E & Halls T D J, *J Indian Chem Soc*, 62 (1985) 916.
- 15 Russell A & Frye J R, *Org Syn Coll*, 3 (1955).